

CONTENT OF VITAMIN E IN PALM OIL AND ITS ANTIOXIDANT ACTIVITY*


AUGUST 1994

28

by: AB GAPOR MD TOP

PORIM INFORMATION SERIES

ISSN 0128-5726



The occurrence of an anti-sterility factor which was characterized as a vitamin was first reported in 1922; later in 1925, it was given the designation vitamin E. It is now known that vitamin E comprises two groups of related compounds, known as tocopherols and tocotrienols and that, in nature, there are four homologues in each group (*Figure 1*).

Vitamin E is a fat-soluble vitamin which is present in small amount in vegetable oils. Studies on the identity of vitamin E compounds in oils and fats have shown that by contrast with the situation in other vegetable oils, vitamin E in palm oil mainly composed of tocotrienols (*Table 1*).

In view of the fact that most vegetable oils including palm oil are consumed in processed forms, we have carried out studies on the vitamin E content of crude and processed palm oil products and the results are given in *Table 2*. The vitamin E content in processed oils was slightly lower than in the corresponding crude oils. It was also found that most of the vitamin E lost during processing went into the palm fatty acid distillate during the step of deacidification/deodorization.

In order to have a better understanding of the role of vitamin E in palm oil, the antioxidant activities of its major components have been studied: α -tocotrienol, γ -tocotrienol and δ -tocotrienol were added individually to vitamin E-free RBD palm olein at various concentration from 200 to 2000 ppm and the

stability of the samples, expressed as the induction period (IP), was measured by using the Rancimat 617 apparatus. The results as given in *Figure 2* indicated that the order of antioxidant activity of individual tocotrienols is as follows :

$$\gamma - \text{tocotrienol} \geq \delta > \alpha - \text{tocotrienol}$$

It was found that γ - tocotrienol had about twice the activity of α - tocotrienol and that at 200 ppm, α

TABLE 1. TYPICAL COMPOSITION OF VITAMIN E IN CRUDE PALM OIL

Compound	% Composition
α - tocopherol	22
α - tocotrienol	20
γ - tocotrienol	46
δ - tocotrienol	12

TABLE 2. VITAMIN E CONTENT (ppm) IN CRUDE AND PROCESSED PALM OILS

Sample	Range	Average	n
Crude palm Oil	708-1141	843	33
RBD palm oil	378-890	581	33
Crude palm olein	880-1129	998	12
RBD palm olein	559-902	716	12
Crude palm stearin	426-552	489	2
RBD palm stearin	348-381	365	2


tocotrienol improved the stability of the substrate by a factor of about 6.3. It was also noted that vitamin E-free RBD palm olein had a relatively low stability (IP ca. 2hr) compared with fresh RBD palm olein (IP ca. 48hr).



It is obvious that the vitamin E components in palm oil play an important role in stabilizing it. The presence of vitamin E is also important nutritionally and it has been suggested by many researchers recently that vitamin E, particularly tocotrienols, are


anti-carcinogenic and cholesterol reducing. Further research and development on vitamin E with particular reference to tocotrienols is being actively pursued.

THE TOCOPHEROLS



-Tocopherol	-Tocopherol	R ₁	R ₂	R ₃
α -	5,7,8 - Trimethyl	CH ₃	CH ₃	CH ₃
β -	5,8 - Dimethyl	CH ₃	H	CH ₃
γ -	7,8 - Dimethyl	H	CH ₃	CH ₃
δ -	8 - Methyl	H	H	CH ₃


THE TOCOTRIENOLS



-Tocotrienol	-Tocotrienol	R ₁	R ₂	R ₃
α -	5,7,8 - Trimethyl	CH ₃	CH ₃	CH ₃
β -	5,8 - Dimethyl	CH ₃	H	CH ₃
γ -	7,8 - Dimethyl	H	CH ₃	CH ₃
δ -	8 - Methyl	H	H	CH ₃

Note: α - Tocopherol, α -, γ - and δ - tocotrienols are the major homologues in palm oil.

Figure 1. Naturally Occurring Tocopherols and Tocotrienols.



CONCENTRATION OF TOCOTRIENOLS (ppm)

- Vitamin E - free RBD Palm Olein (substrate) - Control
- Substrate + α - tocotrienol
- Substrate + γ - tocotrienol
- Substrate + δ - tocotrienol

Figure 2. Antioxidant Activities of Tocotrienols in Vitamin E - free RBD Palm Olein.

Printed with Palm Oil - Based ink by PORIM

For more information kindly contact:

Director-General
P. O. Box 10620
50720 Kuala Lumpur
Malaysia

24229
 Pusat Maklumat
 Sawit



1

ITS ANTIOXIDANT ACTIVITY

12126 / 2