

TOCOTRIENOL AND CURCUMIN ENRICHED TRANSDERMAL CREAM

KANGA RANI SELVADURAY; KALANITHI NESARETNAM; PREMDASS RAMDAS; FU JU YEN; PUVANESWARI MEGHANATHAN; GHAZALI ABDUL RAZAK and ONG THEAN HUAT



MPOB INFORMATION SERIES • ISSN 1511-7871 • JUNE 2015

MPOB TT No. 586

A topical fluid or cream formulation wherein the main ingredient is tocotrienol, and curcumin is added to enhance the anti-cancer effects of tocotrienols. The composition is useful for the treatment or prevention of a cancer, a tumour or an inflammatory disorder, particularly breast cancer.

In one embodiment of the present invention is a composition containing an active amount of tocotrienol or derivative thereof as the main ingredient; and an addition of curcumin, for prevention or inhibition of a cancer and/or a tumour in a mammal, preferably breast cancer,

wherein curcumin is added to the composition to enhance the anti-cancer effect in tocotrienol.

It is to be noted that tocotrienols have potent anti-cancer and anti-inflammatory effects and the addition of curcumin enhances these effects in the product (*Table 1*).

THE TECHNOLOGY

- Production of a compound containing a mixture of tocotrienol and curcumin at a certain combination for optimum synergistic effects.
- The cream form of the compound containing the mixture for transdermal application.

TABLE 1. INHIBITORY EFFECTS OF TRF, CURCUMIN, CURCUMIN + 1 μ M TRF, TRF + 1 μ M CURCUMIN ON MCF-7 HUMAN BREAST CANCER CELLS

Inhibition of proliferation, IC ₅₀			
Curcumin	Curcumin + 1 μ M TRF	TRF	TRF + 1 μ M Curcumin
25.3 μ M	25.7 μ M	30.4 μ M	15 μ M


THE PRODUCT



Tocotrienols from palm oil



Curcumin from turmeric



Tocotrienol + Curcumin cream

ISSN 1511-7871



9 771511 787001

Malaysian Palm Oil Board, Ministry of Plantation Industries and Commodities, Malaysia

6 Persiaran Institusi, Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia. Tel: 03-8769 4400

Fax: 03-8925 9446

Website: www.mpob.gov.my



THE BENEFITS

- An application cream containing combination of tocotrienols and curcumin for reducing inflammation and prevention of cancer.
- A 2-in-1 convenient and cost-effective health product.

REFERENCES

LEV-ARI, S; ZINGER, H; KAZANOV, D; YONA, D; BEN-YOSEF, R; STARR, A; FIGER, A and ARBER, N (2005). Curcumin synergistically potentiates the growth inhibitory and pro-apoptotic effects of celecoxib in pancreatic adenocarcinoma cells. *Biomed Pharmacother*, 59 Suppl 2: S276-280.

Malaysian Patent Application No. PI2001000993, patent grant No. 8906960. Synergistic effects of tocotrienols and curcumin.

MCINTYRE, B S; BRISKI, K P; TIRMENSTEIN, M A; FARISS, A M W; GAPOR A and SYLVESTER, P W (2000). Antiproliferative and apoptotic effects of tocopherols and tocotrienols on normal mouse mammary epithelial cells. *Lipids*, 35: 171-180.

MENON, L G; KUTTAN, R and KUTTAN, G (1999). Anti-metastatic activity of curcumin and catechin. *Cancer Lett*, 141(1-2): 159-165.

NESARETNAM, K (2008). Multitargeted therapy of cancer by tocotrienols. *Cancer Lett*, 269: 388-395.

NESARETNAM, K; SELVARAJAN DORASAMY and PHILIPPA DARBRE (2000). Tocotrienols inhibit growth of ZR-75-1 human breast cancer cells. *Int. J. of Food Science. Nutr.* 51: S95-103.

NESARETNAM, K; GUTHRIE, N; CHAMBERS, A F and CAROLL, K K (1995). Effect of tocotrienols on the growth of a human breast cancer cell line in culture. *Lipids*, 30: 1139-1143.

RAMSEWAK, R S; DEWITT, D L and NAIR, M G (2000). Cytotoxicity, antioxidant and anti-inflammatory activities of curcumins I-III from *Curcuma longa*. *Phytomedicine*, 7(4): 303-308.

SEN, C K; KHANNA, S and ROY S (2006). Tocotrienols: vitamin E beyond tocopherols. *Life Sci*, 78: 2098.

For more information, kindly contact:

Director-General
MPOB

6 Persiaran Institusi, Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Tel: 03-8769 4400

Fax: 03-8925 9446

www.mpob.gov.my